Double Longitudinal Spin Asymmetries of Inclusive Charged Pion Production in Polarized p+p Collisions at 200 GeV

Adam Kocoloski (for the STAR Collaboration)

Massachusetts Institute of Technology 77 Massachusetts Avenue Cambridge, MA 02139 USA

A primary goal of the STAR Spin program at RHIC is the measurement of the polarized gluon distribution function ΔG , which can be obtained from a global analysis incorporating measurements of the double spin asymmetry A_{LL} in various final state channels of polarized p+p collisions. Final states with large production cross sections such as inclusive jet and hadron production are analyzed as the program moves towards the measurement of A_{LL} in the theoretically clean channel of prompt photon production. The channels p+p $\rightarrow \pi^{+/-} + X$ are unique in that the ordering of the measurements of A_{LL} in these two channels is sensitive to the sign of ΔG . Moreover, STAR has already established the procedure for the identification of charged pions and the calculation of their production cross-sections over a broad kinematic range.

This contribution will present first measurements of double longitudinal spin asymmetries for inclusive charged pion production extracted from 3 pb⁻¹ of data at $\sqrt{s}=200$ GeV and 50% beam polarizations. The asymmetries are calculated over the transverse momentum region $2 < p_T < 10$ GeV/c and compared with theoretical predictions incorporating several gluon polarization scenarios. A systematic bias introduced by the selection of charged pions from events satisfying electromagnetic energy triggers will be discussed and estimated using Monte Carlo.