

# Accessing the Gluon Polarization through the Double-Helicity Asymmetry in Charged Pion Production at PHENIX

– Astrid Morreale

3 October 2006 Kyoto, Japan

# PIONS



The **PI MESON** plays an important role in our  $\Delta g$  quest:

- **Zero spin** and composed of first generation quarks
- Pseudo scalar under a parity transformation: pion currents **couple** to the **axial vector** current.
- Production of pions proceed from **g-g** and **g-q** initiated sub processes on **proton-proton** collisions

# PARTONIC CONTRIBUTIONS AT MID-RAPIDITY

The pion's fragmentation function contains all long-distance interactions, they are not calculable but they are **universal**:

$$D^{\pi^+}_u > D^{\pi^0}_u > D^{\pi^-}_u, D^{\pi^+}_g = D^{\pi^-}_g$$

q-g starts to dominate for  $P_T > 5 \text{ GeV}$ , pion production in this  $p_T$  range is sensitive to both the gluon and the quark distributions, with different flavours having different weights for each pion species







Figure 2: Relative fractional contributions of partonic processes to mid-rapidity pion production at  $\sqrt{s} = 200 \text{ GeV}$ , calculated by W. Vogelsang.

# Charged Pion $A_{LL}$

Charged Pion  $A_{LL}$  has **sensitivity** to the sign of  $\Delta g$

Just like in the pion's **fragmentation functions** the size of the asymmetries are ordered.

e.g, if  $\Delta g > 0$ , then:  $A^{\pi^+}_{LL} > A^{\pi^0}_{LL} > A^{\pi^-}_{LL}$



**Predictions** for  $A_{LL}$  of  $\pi^+$  and  $\pi^-$  for various  $\Delta g$  scenarios, from M. Stratmann (left and right) and preliminary neutral pion double asymmetries result (center)

## CENTRAL SPECTROMETERS

### Tracking




- Drift Chamber(DC),
- Pad Chambers(PC)

### Electron and Hadron Identification

- Ring Imaging Cerenkov Counter(RICH)
- ElectroMagnetic Calorimeter(EMCal)

### Photons

- ElectroMagnetic Calorimeter (EMCal)



# Double Longitudinal Asymmetries


$$A_{LL}(p_T) = \frac{\sigma_{++} - \sigma_{+-}}{\sigma_{++} + \sigma_{+-}} \rightarrow \frac{1}{P^2} \frac{N_{++}(p_T) - RN_{+-}(p_T)}{N_{++}(p_T) + RN_{+-}(p_T)}, R = \frac{L_{++}}{L_{+-}}$$

$$\sim \frac{\Delta f}{f}$$

Where the  $N_{\text{subscripts}}$  are the observed particle yields when the beams are polarized in different configurations  $P$  is the beam polarization ( $P^2 = P_1 * P_2$ ) and  $R$  is the relative luminosity.

- Average Beam Polarization 49%.
- 0.89 billion triggered events analyzed, equivalent to 2.3 pb<sup>-1</sup>,

# PT SPECTRA OF PARTICLES




Pions start to fire RICH at 4.7 GeV.

PT spectra of positive(left) and negative(right) charged tracks passing detector cuts.

# pT Spectrum Fit to power law.

-Conservative Estimate of Background



<~5% background

Background sources:



- charged hadron tracks randomly associated with a RICH ring.
- conversion electrons with mis-reconstructed momentum (currently no PHENIX tracking close to the beam-pipe.)




# RESULTS

# SINGLE SPIN ASYMETRIES



Parity-violating longitudinal spin asymmetries  
 $A_L$  are expected to be consistent with zero





# Double Spin Asymmetries for Both Charged Species



# Results Compared to Theoretical Parametrizations





# Combined Pion Species Results



Asymmetries for charged pion in Run-05 are compared to neutral pion results. (Run-05  $\pi^0$  left, Run-06 level2 trigger  $\pi^0$  Right)  
\*Left plot includes low  $P_T$  charged pions.


# Run-06 Projections

**Run-06 PHENIX** data with higher statistics and polarization will provide greater sensitivity



Projections for Run-06 for negative pions(left) and positive pions(right)

## FINAL REMARKS



- **Global studies** aid in the understanding of the gluon's contribution to the proton spin.
- Combined analysis of all three pion species at RHIC can provide valuable information on both the sign and **magnitude of Delta-g**
- This analysis will be repeated for **run6 data** as this becomes available

# BACKUP SLIDES






# Spin Structure of the Proton

Quarks and Gluons carry about 50% (each) of the longitudinal momentum

What about *spin*?

Valence quarks(QPM)



~30%

(QCD)



Illustration by Sebastien Pammentier & Astrid Morreale

**Gluons**, sea quarks


~>,=,<0?

Orbital angular momentum contributions



~??


# Hard Scattering Processes in PP collisions



## FACTORIZATION:


It is assumed that a differential cross section can be written as the convolution of a parton density and a hard scattering process

$$\sigma(pp \rightarrow hX) : f_q(x_1) \otimes f_g(x_2) \otimes \hat{\sigma}^{qg \rightarrow q\gamma}(\hat{s}) \otimes D_q^h(z)$$



- Structure functions (experimental input)
- pQCD hard scattering rates (calculable)
- Fragmentation functions (experimental input)

# Polarized Parton Distribution Functions




arXiv:hep-ph/0603213 v2 28 Jun 2006  
M.Hirai, S. Kumano, N.Saito

- Difference in probability between scattering off of a parton with one spin state vs. the other
- Function of  $X_{Bjorken}$ , -momentum fraction of the proton carried by the parton

# CALCULATING $A_{LL}$ -- Yields

After collecting all charged particles that passed our cuts, we then look at the number of particles detected with the **polarized proton beams** in different configurations



There are 120 bunch crossings, each bunch crossing has luminosity information and the spin pattern in each beam.

The charged pion yields are thus collected:

- By Transverse Momentum range (PT Bin)
- By Bunch Crossing.

$A_{LL}$  is Finally calculated by **Fill**.

# Cuts

- $|\text{BBC zvtx}| < \mathbf{30}$  cm
- Track quality **31** or **63**
- $|\text{zed}| < \mathbf{70}$  cm
- $\mathbf{5} < p_T < \mathbf{10}$  GeV/c
- RICH hit ( $\mathbf{n1} > 0$ )
- Shower-shape: prob  $< \mathbf{0.2}$
- $e/\text{momentum} < 0.9$
- $p_T$ -dependent energy cut:
- $\text{emce} > 0.3 + 0.15p_T$
- PC3 matching ( $< \mathbf{3}$ sigma in z and phi)<sup>\*\*\*</sup>
- EMCAL matching ( $< \mathbf{3}$ sigma in z, phi)

# Sources of background

- RICH threshold  $\gamma \sim 35 \rightarrow 0.017, 3.5, 4.7 \text{ GeV}/c$  for e,  $\mu, \pi$
- $e/\pi$  and  $\mu/\pi < 10^{-3}$  for primary e and  $\mu$ 
  - Small!
  - other non-primary background:
    - Conversion electrons generated in front of the DC or inside the X1 layer, which fire the RICH and are often reconstructed with a (false) high momentum
    - Decay or primary charged particles which don't fire the RICH but are randomly associated with a RICH ring. Found random assoc. probability  $\sim 0.4\%$  in d+Au.

# Energy vs. transverse momentum for tracks.





Figure 7: Energy vs. transverse momentum for tracks passing other analysis cuts. Points in black passed the two energy cuts; points in blue did not.

# pT Spectrum for different cuts



Black: No energy cut



**Blue:**  $>1$  GeV

**Magenta:**  $>2$  GeV

**Red:**  $>0.3 + .15 p_T$





# Double Spin Asymmetries vs Fill





Fill by Fill Asymmetries fitted to a constant

PT 5-6 GeV Top



PT 6-7 GeV Middle

PT 7-10 GeV Bottom



Negative pions(Left)  
Positive pions(Right)

# quality


## Quality of the Drift Chamber tracks

- 0 (1) X1 used
- 1 (2) X2 used
- 2 (4) UV found
- 3 (8) UV unique
- 4 (16) PC1 found
- 5 (48) PC1 unique

For example :

$$63 = 48 + 8 + 4 + 2 + 1$$

$$31 = 16 + 8 + 4 + 2 + 1$$



- First look for tracks with X1 and X2 hits
- Remaining unassociated hits go into X1 only and X2 only tracking
- Z coordinates of tracks are defined by PC1-UV-vertex tracking
- When PC choice is unique, stereo wires try to verify. If they do, then UV hits exist and are unique by construction. When they don't, there are no UV hits. If PC has more than one available choice, the UV are consulted for the "best match". In this case one either gets no UV, a best choice, or remaining ambiguity (tied).

## **pc3sdz , pc3sdphi (emcsdz, emcsdphi)**

Difference in  $Z(\text{cm})/\text{phi}(\text{rads})$  between the track model projection and the hit in the pc3 (EMCal) normalized to sigmas. .

## **spc3sdphi**

Difference between the track model projection and the measurement normalized to sigmas, but for the z-swapped projection.

